

9/8/2017 "Life is a journey, not a destination." -Steven Tyler

HW: "Finding Limits Analytically" HW #1, 2, 3, 5

AIM: How do we find limits analytically?

Warm Up:

Finding Limits by Analytic Methods

Observing the graph of a function only can be misleading at times when finding the limit of a function. It is possible to find limits using algebraic techniques and limit theorems.

You will learn to analyze limits by the following methods:

Methods to Analyze Limits:

1. Direct substitution.
2. Principal Limit Theorem
3. Factor, cancellation technique. Then go back to step 1.
4. The conjugate method, rationalize the numerator. Then, go back to step 2.
5. Special trig limits of $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ or $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$
6. L'Hopit ls Rule (presented in Unit 3)

Substitution Theorem

If f is a polynomial function or rational function^(fraction) then $\lim_{x \rightarrow c} f(x) = f(c)$ provided that if f is a rational function the value of the denominator does not equal 0.

EX #1: Find each of the following limits analytically using direct substitution.

A. $\lim_{x \rightarrow 2} (3x^2 - 5x + 4) = 3(2)^2 - 5(2) + 4 = 6$

B. $\lim_{x \rightarrow 2} \frac{x^3 + 1}{x + 1} = \frac{2^3 + 1}{2 + 1} = \frac{9}{3} = 3$

C. $\lim_{x \rightarrow e} \frac{\ln x}{3x} = \frac{\ln e}{3(e)} = \frac{1}{3e}$

$$\ln x \rightarrow \ln_e x$$

$$e^? = x$$

$$\ln e \rightarrow e^? = e$$

$$\ln 5 = x \rightarrow e^x = 5$$

$$D. \lim_{x \rightarrow 4} \sqrt[3]{x+4} = \sqrt[3]{4+4} = \sqrt[3]{8} = 2$$

$$\uparrow = 180^\circ$$

$$E. \lim_{\theta \rightarrow \frac{\pi}{6}} \sin 2\theta = \sin 2(30) = \sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\frac{\pi}{6} = \frac{180}{6} = 30^\circ$$

$$F. \lim_{x \rightarrow 5} \log_3(x+4) = \log_3(5+4) = \log_3 9 = 2$$

Finding Limits of Functions at Undefined Values

Consider the following cases and what happens when you try to evaluate limits by direct substitution.

EX #2: The Factoring or Cancellation Technique

A.	B.	C.
$\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{x^2 - 6x + 8}$ $\frac{2^2 + 2 - 6}{2^2 - 6(2) + 8} = \frac{0}{0}$ <p>indeterminant (hole) point discontinuity</p>	$\lim_{x \rightarrow 4^+} \frac{x^2 + x - 6}{x^2 - 6x + 8}$	$\lim_{x \rightarrow 4^-} \frac{x^2 + x - 6}{x^2 - 6x + 8}$

$$\frac{\text{DOTS}}{x^2 - a^2}$$

$$(x+a)(x-a)$$

Sum/Diff of cubes

$$a^3 + b^3 = (a+b)(a^2 - ab + b^2)$$

$$a^3 - b^3 = (a-b)(a^2 + ab + b^2)$$

S O AP

HW ✓

$$1) \lim_{x \rightarrow 3} \left(\frac{2}{3}x^2 + 3x \right) = \frac{2}{3}(3)^2 + 3(3) = \textcircled{15}$$

$$2) \lim_{t \rightarrow 4} \frac{t-4}{t^2-16} = \frac{1}{4+4} = \textcircled{\frac{1}{8}}$$

$$\frac{\cancel{t-4}}{(\cancel{t-4})(t+4)} = \frac{1}{t+4}$$

$$3) \lim_{x \rightarrow -3} \frac{x^2 - 5x + 6}{2x + 6}$$

$$\lim_{x \rightarrow -3} \text{ is } \textcircled{\text{DNE}}$$

⊗ When we plug in
-3 we get $\frac{30}{0}$

look at the picture
or table of values

$$5) \lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2}$$

⊗ When we plug in 2
we get $\frac{0}{0}$

$$\lim_{x \rightarrow 2} \frac{\cancel{(x-2)}(x^2 + 2x + 4)}{\cancel{x-2}} = \lim_{x \rightarrow 2} x^2 + 2x + 4$$

$$= 2^2 + 2(2) + 4$$


$$= \textcircled{12}$$

Graphically, you can see the limits of the function shown at right. Just because a function is undefined at a value of x doesn't mean that you can't find the limit. Use the graph of the function to determine the value of each limit below.

$$\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{x^2 - 6x + 8} = \underline{-\frac{5}{2}}$$

$$\lim_{x \rightarrow 4^+} \frac{x^2 + x - 6}{x^2 - 6x + 8} = \underline{\infty}$$

$$\lim_{x \rightarrow 4^-} \frac{x^2 + x - 6}{x^2 - 6x + 8} = \underline{-\infty}$$



What is the process for finding discontinuities of a rational function from pre-calculus? (holes, asymptote)

1. Factor everything we can.
Reduce if possible.
2. A Point discontinuity (hole) occurs when the factor that makes the denominator = 0 gets "cancelled".
3. A non-removable discontinuity (asymptote) occurs when the factor that makes the denominator = 0 DOES NOT get cancelled.

You can perform the same algebraic analysis to find the limit of the removable, or point discontinuities and the non-removable, or infinite discontinuities using what we will call the **Factoring or Cancellation Technique**.

$$\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{x^2 - 6x + 8}$$

$$\Rightarrow \frac{(x+3)(x-2)}{(x-4)(x-2)} \Rightarrow \lim_{x \rightarrow 2} \frac{x+3}{x-4}$$

$\frac{2^2 + 2 - 6}{2^2 - 6(2) + 8} = \frac{0}{0}$
 (hole)

asymptote @ $x=4$ hole @ $x=2$

$\lim = \frac{5}{-2}$

Notice the simplified expression above and consider the behavior of this function. Graphically there is a non-removable discontinuity commonly called a

vertical asymptote

at $x = 4$. Because the y -values do not approach one specific value from both sides then the limit does not exist. By using the numerical, graphical, and algebraic techniques together, you can determine the behavior of the simplified function on either side of the vertical asymptote. This is true because the original function and the simplified function agree everywhere except at the

Point discontinuity (hole)

@ $(2, -\frac{5}{2})$.

Determining Behavior of a Function Using One-Sided Limits

$$\lim_{x \rightarrow 4^+} \frac{x^2 + x - 6}{x^2 - 6x + 8}$$

As $x \rightarrow 4^+$

Pick a value: 4.1

Simplified function: $\frac{x+3}{x-4}$

Plug in
 4.1

$$\frac{7.1}{.1} = \frac{+}{+} = +$$



$$\lim_{x \rightarrow 4^+} f(x) = \infty$$

$$\lim_{x \rightarrow 4^-} \frac{x^2 + x - 6}{x^2 - 6x + 8}$$

As $x \rightarrow 4^-$

Pick a value: 3.9

Simplified function: $\frac{x+3}{x-4}$

Plug in
 3.9

$$\frac{6.9}{-.1} = \frac{+}{-} = -$$

$$\lim_{x \rightarrow 4^-} f(x) = -\infty$$

$$\lim_{x \rightarrow 4} f(x) = \text{DNE}$$

HW packet #10, 12

10) Plug in 2 get $\frac{28}{0}$ ~~DNE~~ ∞ $\lim_{x \rightarrow 2^+}$ Plug in # slightly to the right of 2 $-\infty$

Try 2.1

$$\frac{3(2.1)^2 + 7(2.1) + 2}{(2.1)^2 - 4} = \frac{29.93}{0.41}$$

Positive

The Limit is ∞ 12) Plug in $\frac{3}{2}$ get $\frac{0}{0}$ Test
Tues 9/26

$$\frac{8x^3 - 27}{2x - 3} = \frac{\cancel{(2x-3)}((2x)^2 + (2x)(3) + 3^2)}{\cancel{2x-3}}$$

 $\lim_{x \rightarrow \frac{3}{2}}$ $4x^2 + 6x + 9$ $= 27$

EX #3: The Rationalization Technique or Conjugate Method


A. The graph of $g(x) = \frac{\sqrt{x+2} - 1}{x+1}$ is shown below.

The technique of rationalization can be used to find

$$\lim_{x \rightarrow -1} \frac{\sqrt{x+2} - 1}{x+1}$$

Plug in -1 get $\frac{0}{0}$

⊛ Try Rewrite
conjugate



$$\frac{\sqrt{x+2} - 1}{x+1} \cdot \frac{\sqrt{x+2} + 1}{\sqrt{x+2} + 1} = \frac{x+2-1}{(x+1)(\sqrt{x+2}-1)} = \frac{\cancel{x+1}}{(\cancel{x+1})(\sqrt{x+2}+1)}$$

$$\lim_{x \rightarrow -1} \frac{1}{\sqrt{x+2}+1} = \frac{1}{\sqrt{-1+2}+1} = \left(\frac{1}{2} \right)$$

B. $\lim_{x \rightarrow 5} \frac{x-5}{3 - \sqrt{x+4}}$

EX #4: Find each of the following limits analytically. Show your algebraic steps.

A. $\lim_{x \rightarrow -2} x^3 + 3x^2 - 4x + 5$

B. $\lim_{x \rightarrow \frac{3}{2}} 2x^2 (2x + 3)$

C. $\lim_{x \rightarrow 3} (5x + 1)^{\frac{2}{3}}$

D. $\lim_{x \rightarrow -4} \frac{2x^2 + 7x - 4}{x^2 - x - 20}$

E. $\lim_{x \rightarrow 3} \frac{\sqrt{x+1}-2}{x-3}$

F. $\lim_{x \rightarrow 0} \frac{\sqrt{x+4}-2}{x}$

G. $\lim_{x \rightarrow 0} \frac{\frac{1}{3} - \frac{1}{x+3}}{\frac{x}{1 \cdot 3(x+3)}}$ LCD = $3(x+3)$

$$\Rightarrow \frac{x+3-3}{3x(x+3)} = \frac{\cancel{x}+3-3}{3\cancel{x}(x+3)}$$

Plug in 0 \rightarrow get $\frac{0}{0}$

$$\lim_{x \rightarrow 0} \frac{1}{3(x+3)} = \frac{1}{3(3)} = \left(\frac{1}{9}\right)$$

H. $\lim_{x \rightarrow 2^+} \frac{3x^2 - 7x + 2}{x^2 - 4}$

HW: # 7, 8, 11, 16^{*}
(Try multiplying out
the numerator)